arXiv:1203.0207v1 [hep-ph] 1 Mar 2012

Relativistic Correction to Color Octet J/ψ Production at Hadron Colliders

Guang-Zhi Xu $^{(a,b)}$,[*] Yi-Jie Li $^{(a,b)}$,[†] Kui-Yong Liu $^{(b)}$,[‡] and Yu-Jie Zhang $^{(a)}$[§]
(a) Key Laboratory of Micro-nano Measurement-Manipulation and Physics (Ministry of Education) and School of Physics, Beihang University, Beijing 100191, China
(b) Department of Physics, Liaoning University, Shenyang 110036 , China

(Dated: February 24, 2013)

The relativistic corrections to the color-octet J/ψ hadroproduction at the Tevatron and LHC are calculated up to $\mathcal{O}(v^2)$ in nonrelativistic QCD factorization frame. The short distance coefficients are obtained by matching full QCD with NRQCD results for the subprocess $g+g\to J/\psi({}^1S_0^{[8]}, {}^3S_1^{[8]}, {}^3P_J^{[8]})+g$. The short distance coefficient ratios of relativistic correction to LO for CO states ${}^1S_0^{[8]}$, ${}^3S_1^{[8]}$ and ${}^3P_J^{[8]}$ at large p_T are approximately -5/6, -11/6, and -31/30 respectively, and it is 1/6 for color singlet state ${}^3S_1^{[1]}$. If the long distance matrix elements are estimated through velocity scaling rule with adopting $v^2=0.23$, the LO cross sections of color-octet states are reduced by about a factor of 20 40% at large p_T at both Tevatron and LHC. These results indicate that relativistic corrections may play an important role in J/ψ production and polarization at the Tevatron and LHC.

PACS numbers: 12.38.Bx,12.39.St,13.85.Ni,14.40.Gx

I. INTRODUCTION

Heavy quarkonium is an excellent candidate to probe QCD from the high energy to the low energy regimes. Nonrelativistic QCD (NRQCD) factorization formalism was established[1] to describe the production and decay of heavy quarkonium. In the NRQCD approach, the production and decay of heavy quarkonium is factored to short distance coefficients part and long distance matrix elements(LDMEs) part. The short distance coefficients indicate the creation or annihilation of a heavy quark pair and can be calculated perturbatively with the expansions by the strong coupling constant α_s. However, the LDMES, which represented the evolvement of a free heavy quark pair into a bound state, can be scaled by the relative velocity between the quark and antiquark v and obtained by lattice QCD or extracted from the experiment. The color-octet mechanism (COM) was introduced here. The heavy quark pair should be a color-singlet (CS) bound state at long distances, but it may be in a color-octet (CO) state at short distances. NRQCD had achieved great success since it was proposed. The COM was applied to cancel the infrared divergences in the decay widths of P-wave [2, 3] and D-wave[4, 5] heavy quarkonium. However, difficulties were still encountered. The large discrepancy between the experimental data and the theoretical calculation of J/ψ and ψ' unpolarization and polarization production at Tevatron is a very interesting puzzle that can verify NRQCD when solved[6–8]. Theoretical prediction with COM contributions was introduced fitted well with the experimental data of the J/ψ production at the Tevatron[9]. However, the CO contributions from gluon fragmentation indicated that the J/ψ was transversely polarized at large p_T, inconsistent with the experimental data[10].

The NLO QCD corrections of CS J/ψ hadroproduction was calculated to resolve the puzzle of J/ψ hadroproduction[11, 12]. The calculation enhanced the cross sections at large p_T by approximately an order of magnitude. However, the large discrepancy between the CS predictions and the experimental data remained unsolved. The relativistic correction to CS J/ψ hadroproduction were very small[13]. At the same time, the NLO QCD corrections of COM J/ψ hadroproduction were calculated, which eventually provided a possible solution to the long-standing J/ψ polarization puzzle [14, 15]. A similar large discrepancy was found in the double charmonium production at B factories[16–18]. Much work had been performed, and apparently these discrepancies could be resolved by including NLO QCD corrections[19–22] and relativistic corrections[23–26]. The data from B factories pointed out that the COM LDMEs of J/ψ production may be smaller than previously expected[25–29]. (More information can

---

[*] Electronic address: still200@gmail.com
[†] Electronic address: yijiegood@gmail.com
[‡] Electronic address: liukuiyong@lnu.edu.cn
[§] Electronic address: nophy0@gmail.com

be found in Refs.[30–32]).

In this paper, the effect of relativistic corrections to the COM J/ψ hadroproduction at Tevatron and LHC were estimated based on NRQCD. The short distance coefficients were calculated up to $\mathcal{O}(v^2)$. Many free LDMEs were realized at $\mathcal{O}(v^2)$, which were estimated according to the velocity scaling rules within NRQCD[33].

The paper is organized as follows. In Section II, the frame of calculation is introduced for the relativistic correction of both the S- and P-wave states in NRQCD frame. Section III provides the numerical result. Finally, a brief summary of this work is presented.

II. RELATIVISTIC CORRECTIONS OF CROSS SECTION IN NRQCD

The inclusive production of J/ψ was dominant by the gluon-gluon fusion process at high energy hadron colliders. We can compute the cross section at the level of parton. Finally, the cross section of the hadronic level is obtained by integrating the product of the partonic cross section and the gluon distribution function over momentum fraction.

$$g(k_1) + g(k_2) \rightarrow J/\psi(P) + g(k_3).$$

As the beginning of this paper, in NRQCD, the calculation to production of charmonia is divided in two part: short distance coefficients and LDMEs. So the above process can be written as:

$$\frac{d\sigma}{dt}(g + g \to J/\psi + g) = \sum_n \frac{F_n}{m_c^{d_n-4}} \langle \mathcal{O}_n^{J/\psi} \rangle. \tag{1}$$

On the right-hand side of the equation, the cross section is expanded to sensible Fock states noted by the subscript n. F_n, i.e., short distance coefficients, which describe the process that produces intermediate $Q\bar{Q}$ in a short range before heavy quark and antiquark hadronization to the physical meson state. $\langle \mathcal{O}_n^{J/\psi} \rangle$ are the long distance matrix elements that represent the hadronization $Q\bar{Q}$ evolutes to the CS final state by emitting soft gluons. $\mathcal{O}_n^{J/\psi}$ are local four fermion operators. The factor of $m_c^{d_n-4}$ is introduced to make F_n dimensionless.

In this section, our calculation on the differential cross section for this process in the NRQCD factorization formula is divided into two parts, namely, kinematics and short distance coefficients.

A. Kinematics

We denote the three relative momenta between heavy quark and antiquark as $2\vec{q}$, with $|\vec{q}| \sim m_c v$, in J/ψ rest frame, where m_c are the mass of charm quark and v is the three relative velocities between quark and antiquark. Thus, the momenta for the quark and antiquark are expressed as[24, 34, 35]

$$\begin{aligned} p_c &= (E_q, \vec{q}), \\ p_{\bar{c}} &= (E_q, -\vec{q}). \end{aligned} \tag{2}$$

where $E_q = \sqrt{m_c^2 + |\vec{q}|^2}$ is the rest energy of both the quark and antiquark, and $2E_q$ is the invariable mass of J/ψ. When boosting to an arbitrary frame,

$$p_c \to \tfrac{1}{2}P + q, \quad p_{\bar{c}} \to \tfrac{1}{2}P - q. \tag{3}$$

where P is the four momenta of J/ψ, and q receives the boost from $(0, \vec{q})$.

The Lorentz invariant Mandelstam variables are defined as

$$s = (k_1 + k_2)^2 = (P + k_3)^2,$$

$$t = (k_1 - P)^2 = (k_2 - k_3)^2,$$

$$u = (k_1 - k_3)^2 = (k_2 - P)^2.$$

with the relationship $s+t+u=P^2=4E_q^2$. Here, when expand the amplitude with $|\vec{q}|$ up to $O(|\vec{q}|^2)$, great care must be taken to ensure that that t and u are dependent on $|\vec{q}|$. Another two natural Lorentz invariant variables are the t' and u' corresponding to t and u when the effect of $|\vec{q}|$ is considered. One can easily get that they satisfy

$$t=\frac{s-4E_q^2}{s-4m_c^2}t', \qquad u=\frac{s-4E_q^2}{s-4m_c^2}u'. \tag{4}$$

In our subsequent calculation, we adopt $t(u)$ to represent $t'(u')$ directly for simplification.

The `FeynArts` [36] package was used to generate Feynman diagrams and amplitudes, and the `FeynCalc` [37] package was used to handle amplitudes. The numerical phase space was integrated with Fortran.

B. Short-distance coefficients calculation

According to NRQCD factorization, the differential cross section of gluon-gluon fusion up to next order in v^2 to CS state ${}^3S_1^{[1]}$ and each CO states ${}^1S_0^{[8]}$, ${}^3S_1^{[8]}$, ${}^3P_J^{[8]}$, can be expressed as

$$\begin{aligned} d\sigma &= d\sigma_{lo}[{}^3S_1^{[1]}]+d\sigma_{lo}[{}^1S_0^{[8]}]+d\sigma_{lo}[{}^3S_1^{[8]}]+d\sigma_{lo}[{}^3P_J^{[8]}] \\ &+ d\sigma_{rc}[{}^3S_1^{[1]}]+d\sigma_{rc}[{}^1S_0^{[8]}]+d\sigma_{rc}[{}^3S_1^{[8]}]+d\sigma_{rc}[{}^3P_J^{[8]}]. \end{aligned} \tag{5}$$

In this expression, relativistic correction parts, denoted as 'rc', can easily be distinguished from LO, denoted as 'lo'. In addition, each differential cross section to different Fock states should be divided in short distance coefficient part and LDMEs. We can introduce $F({}^{2s+1}L_J)$ to express the short distance coefficient of the LO cross section, corresponding to $G({}^{2s+1}L_J)$ for relativistic correction. Many LDMEs are present, all of which are denoted by $\langle\mathcal{O}^{J/\psi}({}^{2s+1}L_J)\rangle$ and $\langle\mathcal{P}^{J/\psi}({}^{2s+1}L_J)\rangle$ for the LO and relativistic correction term respectively. The explicit expressions of the eight four-fermion operators are[1]:

$$\begin{aligned}
<0|\mathcal{O}^{J/\psi}({}^3S_1^{[1]})|0> &= <0|\chi^\dagger\sigma^i\psi(a_\psi^\dagger a_\psi)\psi^\dagger\sigma^i\chi|0>, \\
<0|\mathcal{P}^{J/\psi}({}^3S_1^{[1]})|0> &= <0|\frac{1}{2}[\chi^\dagger\sigma^i\psi(a_\psi^\dagger a_\psi)\psi^\dagger\sigma^i(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\chi+h.c.]|0>, \\
<0|\mathcal{O}^{J/\psi}({}^1S_0^{[8]})|0> &= <0|\chi^\dagger T^a\psi(a_\psi^\dagger a_\psi)\psi^\dagger T^a\chi|0>, \\
<0|\mathcal{P}^{J/\psi}({}^1S_0^{[8]})|0> &= <0|\frac{1}{2}[\chi^\dagger T^a\psi(a_\psi^\dagger a_\psi)\psi^\dagger T^a(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\chi+h.c.]|0>, \\
<0|\mathcal{O}^{J/\psi}({}^3S_1^{[8]})|0> &= <0|\chi^\dagger T^a\sigma^i\psi(a_\psi^\dagger a_\psi)\psi^\dagger T^a\sigma^i\chi|0>, \\
<0|\mathcal{P}^{J/\psi}({}^3S_1^{[8]})|0> &= <0|\frac{1}{2}[\chi^\dagger T^a\sigma^i\psi(a_\psi^\dagger a_\psi)\psi^\dagger T^a\sigma^i(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2\chi+h.c.]|0>, \\
<0|\mathcal{O}^{J/\psi}({}^3P_J^{[8]})|0> &= <0|\chi^\dagger T^a(-\frac{i}{2}\overleftrightarrow{D^i}\sigma^j)\psi(a_\psi^\dagger a_\psi)\psi^\dagger T^a(-\frac{i}{2}\overleftrightarrow{D^i}\sigma^j)\chi|0>, \\
<0|\mathcal{P}^{J/\psi}({}^3P_J^{[8]})|0> &= <0|\frac{1}{2}[\chi^\dagger T^a(-\frac{i}{2}\overleftrightarrow{D^i}\sigma^j)\psi(a_\psi^\dagger a_\psi)\psi^\dagger T^a(-\frac{i}{2}\overleftrightarrow{\mathbf{D}})^2(-\frac{i}{2}\overleftrightarrow{D^i}\sigma^j)\chi+h.c.]|0>.
\end{aligned} \tag{6}$$

Where χ and ψ are the Pauli spinors describing anti-charm quark creation and charm quark annihilation respectively. σ is the Pauli matrices and $\mathbf{D}$ is the gauge-covariant derivative with $\overleftrightarrow{\mathbf{D}}=\overrightarrow{\mathbf{D}}-\overleftarrow{\mathbf{D}}$. T is the $SU(3)$ color matrix.It should be noted that

$$<0|\mathcal{O}^{J/\psi}({}^3P_J^{[8]})|0>=(2J+1)(1+\mathcal{O}(v^2))<0|\mathcal{O}^{J/\psi}({}^3P_0^{[8]})|0>. \tag{7}$$

So there are 6 LDMEs for P-wave and 6 LDMEs for S-wave. The long distance matrix elements of heavy quarkonium decay may be determined by potential model[23, 39] or lattice calculations[40], and by phenomenological extraction from experimental data[13, 41]. But it's very difficult for production LDMEs. To simplify the discussion of the numerical result, it is assumed that

$$\begin{aligned} <0|\mathcal{O}^{J/\psi}({}^3P_J^{[8]})|0> &= (2J+1)<0|\mathcal{O}^{J/\psi}({}^3P_0^{[8]})|0>, \\ <0|\mathcal{P}^{J/\psi}({}^3P_J^{[8]})|0> &= (2J+1)<0|\mathcal{P}^{J/\psi}({}^3P_0^{[8]})|0>. \end{aligned} \tag{8}$$

For simplicity, we can estimate the relation between their order as[38, 39].

$$|\vec{q}|^2 = \frac{\langle \mathcal{P}^{J/\psi(^{2s+1}L_J)}\rangle}{\langle \mathcal{O}^{J/\psi(^{2s+1}L_J)}\rangle}[1+\mathcal{O}(v^4)]. \tag{9}$$

$|\vec{q}|$ has defined before and equal to $m_c v$. For charmonia, $v^2 \sim 0.25$. Thus the total cross section can be divided in four sub cross section which noted by each Fock state to be calculated respectively.

The short distance coefficients can be evaluated by matching the computations of perturbative QCD and NRQCD:

$$d\sigma\Big|_{pert\ QCD} = \sum_n \frac{F_n}{m_c^{d_n-4}}\langle \mathcal{O}_n^{c\bar{c}}\rangle\Big|_{pert\ NRQCD}. \tag{10}$$

The covariant projection operator method should be adopted to compute the expression on the left hand of the equation. Using this method, spin-singlet and spin-triplet combinations of spinor bilinears in the amplitudes can be written in covariant form. For the spin-singlet case:

$$\begin{aligned}&\sum_{s\bar{s}} v(s)\bar{u}(\bar{s})\langle \frac{1}{2}, s; \frac{1}{2}, \bar{s}|0,0\rangle \\ &= \frac{1}{2\sqrt{2}(E_q+m)}(-\not{p}_{\bar{c}}+m_c)\gamma_5\frac{\not{P}+2E_q}{2E_q}(\not{p}_c+m_c).\end{aligned} \tag{11}$$

For spin-triplet case, the expression is defined as:

$$\begin{aligned}&\sum_{s\bar{s}} v(s)\bar{u}(\bar{s})\langle \frac{1}{2}, s; \frac{1}{2}, \bar{s}|1,S_z\rangle \\ &= \frac{1}{2\sqrt{2}(E_q+m)}(-\not{p}_{\bar{c}}+m_c)\not{\epsilon}\frac{\not{P}+2E_q}{2E_q}(\not{p}_c+m_c).\end{aligned} \tag{12}$$

Where ϵ denotes the polarization vector of the spin-triplet state. In our calculation, Dirac spinors are normalized as $\bar{u}u = -\bar{v}v = 2m_c$.

The differential cross section of each state then satisfies:

$$d\sigma(^{(2s+1)}L_J)\sim\overline{\sum}|\mathcal{M}(g+g\to(c\bar{c})(^{(2s+1)}L_J)+g)|^2\langle \mathcal{O}^{J/\psi}(^{2s+1}L_J)\rangle. \tag{13}$$

Where the symbol $\overline{\sum}$ is the sum of the final state color and polarization as well as the average initial states. According to this expression and Eq.(6), expanding the cross section to $\mathcal{O}(v^2)$ is to expand the amplitude squared on the right side of the above expression to $\mathcal{O}(|\vec{q}|^2)$.

Next, we prepare to expand the short distance coefficients to the next order in v^2. First, we expand each Fock state amplitude, including the S-wave and P-wave states, in term of the relative momentum $|\vec{q}|$:

$$\begin{aligned}&\mathcal{M}(g+g\to(c\bar{c})(^3S_1^{[1,8]})+g) \\ &= \epsilon_\rho(\mathcal{M}_t^\rho\Big|_{q=0} + \frac{1}{2}q^\alpha q^\beta\frac{\partial^2(\sqrt{\frac{m_c}{E_q}}\mathcal{M}_t^\rho)}{\partial q^\alpha\partial q^\beta}\Big|_{q=0}) + \mathcal{O}(q^4).\end{aligned} \tag{14}$$

$$\begin{aligned}&\mathcal{M}(g+g\to(c\bar{c})(^1S_0^{[8]})+g) \\ &= \mathcal{M}_s\Big|_{q=0} + \frac{1}{2}q^\alpha q^\beta\frac{\partial^2(\sqrt{\frac{m_c}{E_q}}\mathcal{M}_s)}{\partial q^\alpha\partial q^\beta}\Big|_{q=0} + \mathcal{O}(q^4).\end{aligned} \tag{15}$$

$$\begin{aligned}&\mathcal{M}(g+g\to(c\bar{c})(^3P_J^{[8]})+g) = \epsilon_\rho(s_z)\epsilon_\sigma(L_z)(\frac{\partial\mathcal{M}_t^\rho}{\partial q^\sigma}\Big|_{q=0} \\ &+\frac{1}{2}q^\alpha q^\beta\frac{\partial^3(\sqrt{\frac{m_c}{E_q}}\mathcal{M}_t^\rho)}{\partial q^\alpha\partial q^\beta\partial q^\sigma}\Big|_{q=0}) + \mathcal{O}(q^4).\end{aligned} \tag{16}$$

The factor $\sqrt{\frac{m_c}{E_q}}$ comes from the relativistic normalization of $c\bar{c}$ state. Odd power terms of four-momentum q vanish in either the S-wave or the P-wave amplitudes, where $\mathcal{M}_t$ and $\mathcal{M}_s$ are inclusive production amplitudes to triplet and singlet $c\bar{c}$ respectively.

$$\mathcal{M}_s = \sum_{s\bar{s}} \sum_{ij} \langle \frac{1}{2}, s; \frac{1}{2}, \bar{s} | 0, 0 \rangle \langle 3i; \bar{3}j | 1, 8a \rangle \mathcal{A}(g + g \to c^i + \bar{c}^j).$$

$$\mathcal{M}_t = \sum_{s\bar{s}} \sum_{ij} \langle \frac{1}{2}, s; \frac{1}{2}, \bar{s} | 1, S_z \rangle \langle 3i; \bar{3}j | 1, 8a \rangle \mathcal{A}(g + g \to c^i + \bar{c}^j).$$

In the process of expanding the amplitude squared in terms of $|\vec{q}|$, it needs to be integrated over the space angle to four-momentum q. We can obtain the following formulas:

$$\int d\Omega Y_{00}^*(\hat{q}) Y_{00}(\hat{q}) q^\alpha q^\beta = \frac{1}{3} |\vec{q}|^2 \Pi^{\alpha\beta}, \tag{17}$$

$$\sum_{L_z} \int d\Omega Y_{1L_z}^*(\hat{q}) Y_{1L_z}(\hat{q}) q^\alpha q^\beta = \frac{1}{3} |\vec{q}|^2 \Pi^{\alpha\beta}, \tag{18}$$

$$\begin{aligned} &\sum_{L_z} \int d\Omega Y_{1L_z}^*(\hat{q}) Y_{1L_z}(\hat{q}) q^\alpha q^\beta q^\sigma q^\rho \\ &= \frac{1}{5} |\vec{q}|^4 \left(\Pi^{\alpha\beta} \Pi^{\sigma\rho} + \Pi^{\alpha\sigma} \Pi^{\beta\rho} + \Pi^{\alpha\rho} \Pi^{\beta\sigma} \right). \end{aligned} \tag{19}$$

where $\Pi^{\alpha\beta} = -g^{\mu\nu} + \frac{P^\mu P^\nu}{P^2}$.

Subsequently, the amplitude squared up to the next order can be obtained

$$\sum |\mathcal{M}(^3S_1^{[1,8]})|^2 = \mathcal{M}_t^\rho(0) \mathcal{M}_t^{\lambda *}(0) \sum \epsilon_\rho \epsilon_\lambda^* + \frac{1}{3} |\vec{q}|^2 \left[(\Pi^{\alpha\beta} \frac{\partial^2 (\sqrt{\frac{m_c}{E_q}} \mathcal{M}_t^\rho)}{\partial q^\alpha \partial q^\beta}) \Big|_{q=0} \mathcal{M}_t^{\lambda *}(0) \right] \sum \epsilon_\rho \epsilon_\lambda^* + \mathcal{O}(v^4). \tag{20}$$

$$\sum |\mathcal{M}(^1S_0^{[8]})|^2 = \mathcal{M}_s(0) \mathcal{M}_s^*(0) + \frac{1}{3} |\vec{q}|^2 \left[(\Pi^{\alpha\beta} \frac{\partial^2 (\sqrt{\frac{m_c}{E_q}} \mathcal{M}_s)}{\partial q^\alpha \partial q^\beta}) \Big|_{q=0} \mathcal{M}_s^*(0) \right] + \mathcal{O}(v^4). \tag{21}$$

$$\begin{aligned} \sum |\mathcal{M}(^3P_J^{[8]})|^2 &= |\vec{q}|^2 \Pi^{\alpha\beta} \frac{\partial \mathcal{M}_t^\rho}{\partial q^\alpha} \Big|_{q=0} \frac{\partial \mathcal{M}_t^{\lambda *}}{\partial q^\beta} \Big|_{q=0} \sum \epsilon_\rho \epsilon_\lambda^* \\ &+ \frac{1}{15} |\vec{q}|^4 \left[(\Pi^{\alpha\beta} \Pi^{\sigma\tau} (\frac{\partial^3}{\partial q^\alpha \partial q^\sigma \partial q^\tau} + \frac{\partial^3}{\partial q^\sigma \partial q^\alpha \partial q^\tau} + \frac{\partial^3}{\partial q^\tau \partial q^\sigma \partial q^\alpha}) (\sqrt{\frac{m_c}{E_q}} \mathcal{M}_t^\rho)) \frac{\partial \mathcal{M}_t^{\lambda *}}{\partial q^\beta} (\sum \epsilon_\rho \epsilon_\lambda^*) \right]_{q=0} \\ &+ \mathcal{O}(v^6). \end{aligned} \tag{22}$$

Any term, which is in the order of $|\vec{q}|^2$, must not be missed to obtain the correction up to the order of v^2. In the three expressions above, each first term on the right side of the equation can be expressed in terms of kinematics variables s, t, u. Here t, u depends on $|\vec{q}|$. The sum of terms in the order of $|\vec{q}|^2$ in the first term as well as all the values of the second term is the result of the next leading order. Polarization sum $\sum \epsilon_\rho \epsilon_\lambda^*$ is equal to $\Pi^{\rho\lambda}$. It should be noted here that the polarization vector ϵ_ρ also depends on q, but the dependence only appears even powers of four momentum q, and the explicit expressions can be found in Ref. [42]. So ϵ_ρ can be carried out in Eq.(20).

Therefore the differential cross section in Eq.(5) takes the following form

$$\begin{aligned}\frac{d\sigma}{dt}(g+g\to J/\psi+g) \;=\;& \left(\frac{F(^3S_1^{[1]})}{m_c^2}\langle 0|\mathcal{O}^{J/\psi}(^3S_1^{[1]})|0\rangle+\frac{G(^3S_1^{[1]})}{m_c^4}\langle 0|\mathcal{P}^{J/\psi}(^3S_1^{[1]})|0\rangle+\right.\\ & \frac{F(^1S_0^{[8]})}{m_c^2}\langle 0|\mathcal{O}^{J/\psi}(^1S_0^{[8]})|0\rangle+\frac{G(^1S_0^{[8]})}{m_c^4}\langle 0|\mathcal{P}^{J/\psi}(^1S_0^{[8]})|0\rangle+\\ & \frac{F(^3S_1^{[8]})}{m_c^2}\langle 0|\mathcal{O}^{J/\psi}(^3S_1^{[8]})|0\rangle+\frac{G(^3S_1^{[8]})}{m_c^4}\langle 0|\mathcal{P}^{J/\psi}(^3S_1^{[8]})|0\rangle+\\ & \left.\frac{F(^3P_0^{[8]})}{m_c^2}\langle 0|\mathcal{O}^{J/\psi}(^3P_0^{[8]})|0\rangle+\frac{G(^3P_0^{[8]})}{m_c^4}\langle 0|\mathcal{P}^{J/\psi}(^3P_0^{[8]})|0\rangle\right)\times\\ & \Big(1+\mathcal{O}(v^4)\Big). \end{aligned}\tag{23}$$

The explicit expressions of the short distance coefficient to the relativistic correction of CO states $^1S_0^{[8]}$ and $^3S_1^{[8]}$ are relegated to the Appendix. The short distance coefficients of CO states $^3P_J^{[8]}$ are too long to list in the Appendix. The result of our relativistic correction of $^3S_1^{[1]}$ is consistent with that of Ref.[13] and was not given in this paper.

III. NUMERICAL RESULT AND DISCUSSION

We adopt the gluon distribution function CTEQ6 PDFs[43]. And the charm quark is set as $m_c = 1.5GeV$. The ratios of the CO short distance coefficient between LO F and it's relativistic correction G at the Tevatron with $\sqrt{s} = 1.96GeV$ and at the LHC with $\sqrt{s} = 7GeV$ or $\sqrt{s} = 14GeV$ are presented in Fig.1. The ratios at the Tevatron and at the LHC are very close. In the large p_T limit,

$$-\frac{M^2}{u}\sim -\frac{M^2}{t}<\frac{M^2}{p_T^2}\sim 0 \tag{24}$$

Then we can expand the short distance coefficients with M. The ratios of first order in the expansion are

$$\begin{aligned} R(^3S_1^{[1]})\Big|_{p_T\gg M} &= \frac{G(^3S_1^{[1]})}{F(^3S_1^{[1]})}\Big|_{p_T\gg M} &\sim\; \frac{1}{6}\\ R(^1S_0^{[8]})\Big|_{p_T\gg M} &= \frac{G(^1S_0^{[8]})}{F(^1S_0^{[8]})}\Big|_{p_T\gg M} &\sim\; -\frac{5}{6}\\ R(^3S_1^{[8]})\Big|_{p_T\gg M} &= \frac{G(^3S_1^{[8]})}{F(^3S_1^{[8]})}\Big|_{p_T\gg M} &\sim\; -\frac{11}{6}\\ R(^3P_0^{[8]})\Big|_{p_T\gg M} &= \frac{G(^3P_0^{[8]})}{F(^3P_0^{[8]})}\Big|_{p_T\gg M} &\sim\; -\frac{31}{30} \end{aligned}\tag{25}$$

It is consistent with the curves in Fig.1.

The ratio of $^3S_1^{[8]}$ is approximately $-11/6$. In large p_T limit, the dominate contribution of this subprocess is $g^*\to c\bar{c}(^3S_1^{[8]})$. The propagator of virtual gluon g^* is proportional to $1/E_q^2$. This term offer a factor of -2 to the ratio $R(^3S_1^{[8]})$.

As discussed in Section II, the LDMEs of relativistic correction are depressed by approximately -0.23 to LO. If the long distance matrix elements are estimated through velocity scaling rule with adopting $v^2 = 0.23$, the LO cross sections of CO states are reduced by about a factor of 20 40% at large p_T at both Tevatron and LHC. Compared with the relativistic corrections to CS state, that ratio is approximately 4%. [1]

---

[1] In Ref.[13], the ratio is about 1% for the LDMEs is different:

$$\langle 0|\mathcal{O}^{J/\psi}(^3S_1^{[1]})|0\rangle = 1.64GeV^3,\qquad \langle 0|\mathcal{P}^{J/\psi}(^3S_1^{[1]})|0\rangle = 0.320GeV^5. \tag{26}$$

The QCD corrections of both CO and CS states had been calculated in [14, 15]. The K factor of NLO QCD corrections is very large for ${}^3P_0^{[8]}$ and ${}^3S_1^{[1]}$, and it is about 1 ± 0.2 for ${}^1S_0^{[8]}$ and ${}^3S_1^{[8]}$. If we fit the matrix elements to NLO in α_s and v^2 at large p_T, $\langle 0|\mathcal{O}^{J/\psi}({}^3S_1^{[1]})|0\rangle$ and $\langle 0|\mathcal{O}^{J/\psi}({}^1S_0^{[8]})|0\rangle$ should be enhance by a factor of 70% and 25% respectively by relativistic corrections. But the relativistic corrections for ${}^3P_0^{[8]}$ and ${}^3S_1^{[1]}$ can be ignored at large p_T for it is much smaller than the QCD corrections And it may affect the size of LDMEs too.



FIG. 1: Ratios of $\mathcal{O}(v^2)$ to $\mathcal{O}(v^0)$ results for CO states. The left diagram's result is calculated at Tevatron and the right one is at LHC. They give pretty close values at large p_T.

IV. SUMMARY

In summary, we calculate the relativistic correction terms to CO states for J/ψ production at the Tevatron and at the LHC. The short distance coefficient ratios of relativistic correction to LO for CO states ${}^1S_0^{[8]}$, ${}^3S_1^{[8]}$ and ${}^3P_J^{[8]}$ at large p_T are approximately -5/6, -11/6, and -31/30 respectively, and it is 1/6 for color singlet state ${}^3S_1^{[1]}$. If the long distance matrix elements are estimated through velocity scaling rule with adopting $v^2 = 0.23$, the LO cross sections of CO states are reduced by about a factor of 20 40% at large p_T at both Tevatron and LHC.Compared with the relativistic corrections to CS state, that ratio is approximately 4%.Thus the result may affect the production and polarization of J/ψ and hadronic colliders. And it may affect the size of LDMEs to.

These results indicate that relativistic corrections may play an important role in J/ψ production and polarization at the Tevatron and LHC.

Then

$$\langle 0|\mathcal{P}^{J/\psi}({}^3S_1^{[1]})|0\rangle / \langle 0|\mathcal{O}^{J/\psi}({}^3S_1^{[1]})|0\rangle / m_c^2 = 0.087, \tag{27}$$

which is much smaller than $v^2 \approx 0.23$.

Acknowledgments

The authors would like to thank Professor K.T. Chao, Z.G. He, Y.Q. Ma, H.S. Shao, and K. Wang for useful discussion and the data of NLO QCD corrections. This work was supported by the National Natural Science Foundation of China (No.10805002, No.10875055, No.11075011), the Foundation for the Author of National Excellent Doctoral Disserta- tion of China (Grant No. 2007B18, No. 201020), the Project of Knowledge Innovation Program (PKIP) of Chinese Academy of Sciences, Grant No. KJCX2.YW.W10 and the Education Ministry of Liaoning Province.

V. APPENDIX: SHORT DISTANCE COEFFICIENTS

The short distance coefficients of ${}^1S_0^{[8]}$ were:

$$
\begin{aligned}
\frac{F({}^1S_0^{[8]})}{m_c^2} &= \frac{1}{16\pi s^2}\frac{1}{64}\frac{1}{4}\frac{1}{N_c^2-1}\times \\
&320\Big[M^{12}\left(t^2+tu+u^2\right)+\left(t^2+tu+u^2\right)^4-M^{10}\left(4t^3+7t^2u+7tu^2+4u^3\right) \\
&+M^8\left(8t^4+21t^3u+27t^2u^2+21tu^3+8u^4\right)-M^6\left(10t^5+35t^4u+57t^3u^2+57t^2u^3+35tu^4+10u^5\right) \\
&+M^4\left(8t^6+33t^5u+66t^4u^2+81t^3u^3+66t^2u^4+33tu^5+8u^6\right) \\
&-M^2\left(t^2+tu+u^2\right)^2\left(4t^3+9t^2u+9tu^2+4u^3\right)\Big]\Big/\Big[M\left(M^2-t\right)^2t\left(M^2-u\right)^2\left(M^2-t-u\right)u(t+u)^2\Big].
\end{aligned}
\tag{28}
$$

$$
\begin{aligned}
\frac{G({}^1S_0^{[8]})}{m_c^4} &= \frac{1}{16\pi s^2}\frac{1}{64}\frac{1}{4}\frac{1}{N_c^2-1} \\
&640\Big[12M^{18}\left(t^2+tu+u^2\right)+5tu(t+u)\left(t^2+tu+u^2\right)^4-5M^{16}\left(11t^3+20t^2u+20tu^2+11u^3\right) \\
&+M^{14}\left(95t^4+280t^3u+358t^2u^2+280tu^3+95u^4\right) \\
&-3M^{12}\left(16t^5+95t^4u+175t^3u^2+175t^2u^3+95tu^4+16u^5\right) \\
&-2M^{10}\left(45t^6+72t^5u+21t^4u^2-22t^3u^3+21t^2u^4+72tu^5+45u^6\right) \\
&+M^8\left(198t^7+678t^6u+1141t^5u^2+1345t^4u^3+1345t^3u^4+1141t^2u^5+678tu^6+198u^7\right) \\
&-M^6\left(180t^8+756t^7u+1583t^6u^2+2224t^5u^3+2446t^4u^4+2224t^3u^5+1583t^2u^6+756tu^7+180u^8\right) \\
&+M^4(85t^9+408t^8u+1000t^7u^2+1637t^6u^3+2028t^5u^4+2028t^4u^5+1637t^3u^6+1000t^2u^7 \\
&+408tu^8+85u^9)-M^2\left(t^3+2t^2u+2tu^2+u^3\right)^2\left(17t^4+30t^3u+30t^2u^2+30tu^3+17u^4\right)\Big]\Big/ \\
&\Big[3M^3\left(M^2-t\right)^3t\left(M^2-u\right)^3u(t+u)^3\left(-M^2+t+u\right)\Big]
\end{aligned}
\tag{29}
$$

The short distance coefficients of ${}^3S_1^{[8]}$ were:

$$
\begin{aligned}
\frac{F({}^3S_1^{[8]})}{m_c^2} &= \frac{1}{16\pi s^2}\frac{1}{64}\frac{1}{4}\frac{1}{N_c^2-1}\frac{1}{3} \\
&256\Big[19M^8\left(t^2+tu+u^2\right)+27\left(t^2+tu+u^2\right)^3-M^6\left(65t^3+111t^2u+111tu^2+65u^3\right) \\
&+M^4\left(100t^4+227t^3u+300t^2u^2+227tu^3+100u^4\right) \\
&-27M^2\left(3t^5+8t^4u+13t^3u^2+13t^2u^3+8tu^4+3u^5\right)\Big]\Big/\Big[3M^3\left(M^2-t\right)^2\left(M^2-u\right)^2(t+u)^2\Big]
\end{aligned}
\tag{30}
$$

$$
\begin{aligned}
\frac{G({}^3S_1^{[8]})}{m_c^4} = {} & \frac{1}{16\pi s^2}\frac{1}{64}\frac{1}{4}\frac{1}{N_c^2-1}\frac{1}{3} \\
& (-512)\Big[M^{14}\left(87t^2+22tu+87u^2\right)+M^{12}\left(-14t^3+335t^2u+335tu^2-14u^3\right) \\
& -2M^{10}\left(399t^4+1612t^3u+2020t^2u^2+1612tu^3+399u^4\right) \\
& +M^8\left(2100t^5+8976t^4u+14497t^3u^2+14497t^2u^3+8976tu^4+2100u^5\right) \\
& -M^6\left(2590t^6+12096t^5u+23855t^4u^2+29314t^3u^3+23855t^2u^4+12096tu^5+2590u^6\right) \\
& +M^4\left(1620t^7+8498t^6u+19905t^5u^2+29152t^4u^3+29152t^3u^4+19905t^2u^5+8498tu^6+1620u^7\right) \\
& -27M^2\left(15t^8+104t^7u+295t^6u^2+510t^5u^3+612t^4u^4+510t^3u^5+295t^2u^6+104tu^7+15u^8\right) \\
& +297tu(t+u)\left(t^2+tu+u^2\right)^3\Big]\Big/\Big[9M^5\left(M^2-t\right)^3\left(M^2-u\right)^3(t+u)^3\Big] \qquad (31)
\end{aligned}
$$

---

[1] G.T. Bodwin, E. Braaten, and G.P. Lepage, Phys. Rev. D **51**, 1125 (1995); **55**, 5853(E) (1997).

[2] Han-Wen Huang and Kuang-Ta Chao, Phys. Rev. **D54**, 3065 (1996) ; *ibid.* **D56**, 7472(E) (1997) (hep-ph/9601283); Phys. Rev. **D55**, 244 (1997) (hep-ph/9605362); Phys. Rev. **D54**, 6850 (1996) ; *ibid.* **D56**, 1821(E) (1997) (hep-ph/9606220).

[3] A. Petrelli, M. Cacciari, M. Greco, F. Maltoni and M.L. Mangano, Nucl. Phys. **B514**, 245 (1998) (hep-ph/9707223).

[4] Z. -G. He, Y. Fan and K. -T. Chao, Phys. Rev. Lett. **101**, 112001 (2008) [arXiv:0802.1849 [hep-ph]].
Z. -G. He, Y. Fan and K. -T. Chao, Phys. Rev. D **81**, 074032 (2010) [arXiv:0910.3939 [hep-ph]].

[5] Y. Fan, Z. -G. He, Y. -Q. Ma and K. -T. Chao, Phys. Rev. D **80**, 014001 (2009) [arXiv:0903.4572 [hep-ph]].

[6] **CDF** Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **69** 3704 (1992).

[7] **CDF** Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **79** 572 (1997).

[8] **CDF** Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **79** 578 (1997).

[9] E.Braaten and S.Fleming, Phys. Rev. Lett. **74**, 3327 (1995). (hep-ph/9411365)

[10] **CDF** Collaboration, Phys. Rev. Lett. **99** 132001 (2007).

[11] J. Campbell, F. Maltoni and F. Tramontano, Phys. Rev. Lett. 98, 252002 (2007). P. Artoisenet, J. M. Campbell, F. Maltoni and F. Tramontano, Phys. Rev. Lett. **102**, 142001 (2009) [arXiv:0901.4352 [hep-ph]]. P. Artoisenet, J. M. Campbell, J. P. Lansberg, F. Maltoni and F. Tramontano, Phys. Rev. Lett. **101**, 152001 (2008) [arXiv:0806.3282 [hep-ph]]. P. Artoisenet, J. P. Lansberg and F. Maltoni, Phys. Lett. B **653**, 60 (2007) [hep-ph/0703129 [HEP-PH]]. H. Haberzettl and J. P. Lansberg, Phys. Rev. Lett. **100**, 032006 (2008) [arXiv:0709.3471 [hep-ph]]. Adam K. Leibovich, Phys. Rev. D **56**, 4412 (1997)(hep-ph/9610381).
G. C. Nayak, J. -W. Qiu and G. F. Sterman, Phys. Rev. Lett. **99**, 212001 (2007) [arXiv:0707.2973 [hep-ph]].

[12] B. Gong and J. -X. Wang, Phys. Rev. D **78**, 074011 (2008) [arXiv:0805.2469 [hep-ph]]. B. Gong and J. -X. Wang, Phys. Rev. Lett. **100**, 232001 (2008) [arXiv:0802.3727 [hep-ph]]. B. Gong, R. Li and J. -X. Wang, arXiv:1102.0118 [hep-ph]. B. Gong, J. -X. Wang and H. -F. Zhang, Phys. Rev. D **83**, 114021 (2011) [arXiv:1009.3839 [hep-ph]]. B. Gong, X. Q. Li and J. -X. Wang, Phys. Lett. B **673**, 197 (2009) [Erratum-ibid. **693**, 612 (2010)] [arXiv:0805.4751 [hep-ph]]. H. S. Chung, S. Kim, J. Lee and C. Yu, Phys. Rev. D **83**, 037501 (2011) [arXiv:1012.1954 [hep-ph]]. Z. -G. He, R. Li and J. -X. Wang, Phys. Rev. D **79**, 094003 (2009) [arXiv:0904.2069 [hep-ph]]. Z. -G. He, R. Li and J. -X. Wang, arXiv:0904.1477 [hep-ph]. D. Li, Y. -Q. Ma and K. -T. Chao,

[13] Y. Fan, Y. -Q. Ma and K. -T. Chao, Phys. Rev. D **79**, 114009 (2009) [arXiv:0904.4025 [hep-ph]].

[14] K. -T. Chao, Y. -Q. Ma, H. -S. Shao, K. Wang and Y. -J. Zhang, arXiv:1201.2675 [hep-ph].

[15] Y. -Q. Ma, K. Wang and K. -T. Chao, Phys. Rev. D **84**, 114001 (2011) [arXiv:1012.1030 [hep-ph]]. Y. -Q. Ma, K. Wang and K. -T. Chao, Phys. Rev. Lett. **106**, 042002 (2011) [arXiv:1009.3655 [hep-ph]]. Y. -Q. Ma, K. Wang and K. -T. Chao, Phys. Rev. D **83**, 111503 (2011) [arXiv:1002.3987 [hep-ph]]. M. Butenschoen and B. A. Kniehl, Phys. Rev. D **84**, 051501 (2011) [arXiv:1105.0820 [hep-ph]]. M. Butenschoen and B. A. Kniehl, Phys. Rev. Lett. **106**, 022003 (2011) [arXiv:1009.5662 [hep-ph]]. M. Butenschoen and B. A. Kniehl, arXiv:1201.3862 [hep-ph].

[16] K. Abe *et al.* [BELLE Collaboration], Phys. Rev. Lett. **89**, 142001 (2002).

[17] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. D **72**, 031101 (2005).

[18] E. Braaten and J. Lee, Phys. Rev. D **67**, 054007 (2003) [Erratum-ibid. D **72**, 099901 (2005)]; K. Hagiwara, E. Kou, and C.F. Qiao, Phys. Lett. B570, 39 (2003). K. -Y. Liu, Z. -G. He and K. -T. Chao, Phys. Rev. D **77**, 014002 (2008) [hep-ph/0408141]. K. -Y. Liu, Z. -G. He and K. -T. Chao, Phys. Rev. D **68**, 031501 (2003) [hep-ph/0305084]. K. -Y. Liu, Z. -G. He and K. -T. Chao, Phys. Rev. D **69**, 094027 (2004) [hep-ph/0301218]. K. -Y. Liu, Z. -G. He and K. -T. Chao, Phys. Lett. B **557**, 45 (2003) [hep-ph/0211181].

[19] Y. J. Zhang, Y. J. Gao and K. T. Chao, Phys. Rev. Lett. **96**, 092001 (2006).

[20] Y. J. Zhang and K. T. Chao, Phys. Rev. Lett. **98**, 092003 (2007); B. Gong and J.X. Wang, arXiv:0904.1103.

[21] B. Gong and J. X. Wang, Phys. Rev. D **77**, 054028 (2008); Phys. Rev. Lett. **100**, 181803 (2008); arXiv: 0904.1103.

[22] Y.J. Zhang, Y.Q. Ma, and K.T. Chao, Phys. Rev. D78, 054006 (2008). K. Wang, Y. -Q. Ma and K. -T. Chao, Phys. Rev. D **84**, 034022 (2011) [arXiv:1107.2646 [hep-ph]]. H. -R. Dong, F. Feng and Y. Jia, JHEP **1110**, 141 (2011) [arXiv:1107.4351 [hep-ph]].

[23] G. T. Bodwin, D. Kang and J. Lee,

[24] Z. G. He, Y. Fan and K. T. Chao, Phys. Rev. D **75**, 074011 (2007) [arXiv:hep-ph/0702239].

[25] Y. Jia, Phys. Rev. D **82**, 034017 (2010) [arXiv:0912.5498 [hep-ph]].

[26] Z. -G. He, Y. Fan and K. -T. Chao, Phys. Rev. D **81**, 054036 (2010) [arXiv:0910.3636 [hep-ph]].

[27] Y. -Q. Ma, Y. -J. Zhang and K. -T. Chao, Phys. Rev. Lett. **102**, 162002 (2009) [arXiv:0812.5106 [hep-ph]].

[28] B. Gong and J. -X. Wang, Phys. Rev. Lett. **102**, 162003 (2009) [arXiv:0901.0117 [hep-ph]].

[29] Y. -J. Zhang, Y. -Q. Ma, K. Wang and K. -T. Chao, Phys. Rev. D **81**, 034015 (2010) [arXiv:0911.2166 [hep-ph]].

[30] Michael Krämer, Prog. Part. Nucl. Phys. **47** 141 (2001) (hep-ph/0106120).

[31] J. P. Lansberg, Int. J. Mod. Phys. A **21**, 3857 (2006) [arXiv:hep-ph/0602091].

[32] N. Brambilla, S. Eidelman, B. K. Heltsley, R. Vogt, G. T. Bodwin, E. Eichten, A. D. Frawley and A. B. Meyer *et al.*, Eur. Phys. J. C **71**, 1534 (2011) [arXiv:1010.5827 [hep-ph]].

[33] G.P.Lepage,L.Magnea,C.Nakhleh,U.Magnea and K.Hornbostel, Phys. Rev. D **46**, 4052 (1992).

[34] C. B. Paranavitane, B. H. J. McKellar and J. P. Ma, Phys. Rev. D **61**, 114502 (2000).

[35] Eric Braaten and Yu-Qi Chen, Phys. Rev. D **54**, 3216 (1996).

[36] J. Küblbeck, M. Böhm, A. Denner, Comput. Phys. Commun. **60** (1990) 165; T. Hahn, Comput. Phys. Commun. **140**, 418 (2001).

[37] R. Mertig, M. Böhm, A. Denner, Comput. Phys. Commun. **64** (1991) 345.

[38] G. T. Bodwin and A. Petrelli, Phys. Rev. D **66**, 094011 (2002) [hep-ph/0205210].

[39] G. T. Bodwin, H. S. Chung, D. Kang, J. Lee and C. Yu, Phys. Rev. D **77**, 094017 (2008) [arXiv:0710.0994 [hep-ph]].

[40] G. T. Bodwin, D. K. Sinclair and S. Kim, Phys. Rev. Lett. **77**, 2376 (1996) [hep-lat/9605023]. G. T. Bodwin, D. K. Sinclair and S. Kim, Phys. Rev. D **65**, 054504 (2002) [hep-lat/0107011]. G. T. Bodwin, J. Lee and D. K. Sinclair, Phys. Rev. D **72**, 014009 (2005) [hep-lat/0503032]. G. T. Bodwin, D. K. Sinclair and S. Kim, Phys. Rev. D **65**, 054504 (2002) [hep-lat/0107011].

[41] H. -K. Guo, Y. -Q. Ma and K. -T. Chao, Phys. Rev. D **83**, 114038 (2011) [arXiv:1104.3138 [hep-ph]].

[42] C. -H. Chang, J. -X. Wang and X. -G. Wu, Phys. Rev. D **70**, 114019 (2004) [hep-ph/0409280].

[43] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky and W. K. Tung, JHEP **0207**, 012 (2002) [hep-ph/0201195].